



A Note on Feller's Strong Law of Large Numbers

Yuan Shih Chow; Cun-Hui Zhang

The Annals of Probability, Vol. 14, No. 3. (Jul., 1986), pp. 1088-1094.

Stable URL:

<http://links.jstor.org/sici?sici=0091-1798%28198607%2914%3A3%3C1088%3AANOFSL%3E2.0.CO%3B2-6>

The Annals of Probability is currently published by Institute of Mathematical Statistics.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ims.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

A NOTE ON FELLER'S STRONG LAW OF LARGE NUMBERS

BY YUAN SHIH CHOW¹ AND CUN-HUI ZHANG

Columbia University and SUNY at Stony Brook

Let X_n , $n \geq 1$, be i.i.d. random variables with common distribution function $F(x)$ and γ_n , $n \geq 1$, be a sequence of constants such that γ_n/n is nondecreasing in n . Set $S_n = X_1 + \cdots + X_n$. The main theorem of this paper gives an integral test which determines the infinite limit points of $\{S_n/\gamma_n\}$. This result extends and combines Feller's (1946) strong law of large numbers (SLLN) and the results of Kesten (1970) and Erickson (1973).

1. Introduction. Let X , X_n , $n \geq 1$, be independent identically distributed random variables with common distribution function $F(x)$. And let γ_n , $n \geq 1$, be a sequence of positive constants such that γ_n/n is nondecreasing in n . Then, the sequence $S_n = X_1 + \cdots + X_n$, $n \geq 1$, is called a random walk and the normalized random walk which we shall study is $\{S_n/\gamma_n\}$.

Define

$$\begin{aligned}\gamma(x) &= \gamma_n, & x = n, \quad n \geq 0, \quad \gamma_0 &= 0 \\ &= \gamma_n + (\gamma_{n+1} - \gamma_n)(x - n), & n \leq x < n + 1, \quad n \geq 0.\end{aligned}$$

And let $\gamma^{-1}(\cdot)$ denote the inverse function of $\gamma(\cdot)$. Set $c = EX/(\lim \gamma_n/n)$ if $E|X| < \infty$ and $c = 0$ otherwise. Feller (1946) obtained the following remarkable result

$$(1.1) \quad \begin{aligned}P\{\lim S_n/\gamma_n = c\} &= 1 \quad \text{iff } E\gamma^{-1}(|X|) < \infty \\ &\text{iff } P\{\limsup |S_n/\gamma_n| < \infty\} = 1,\end{aligned}$$

where iff stands for if and only if.

However, as far as $\limsup S_n/\gamma_n$ and $\liminf S_n/\gamma_n$ are concerned, Feller's strong law of large numbers does not cover the case where $E\gamma^{-1}(|X|) = \infty$. Define

$$\begin{aligned}m_+(x) &= \int_0^x P\{X^+ \geq t\} dt, & x^+ &= \max(x, 0), \\ m_-(x) &= \int_0^x P\{X^- \geq t\} dt, & x^- &= \max(-x, 0), \\ J_+(\gamma) &= \int_0^\infty \min(\gamma^{-1}(x), x/m_-(x)) dF(x),\end{aligned}$$

and

$$J_-(\gamma) = \int_0^\infty \min(\gamma^{-1}(x), x/m_+(x)) d(1 - F(-x)).$$

The following theorem gives an integral test which extends (1.1).

Received September 1984; revised December 1984.

¹Supported by NSF Grant MCS 82-01723.

AMS 1980 subject classifications. Primary 60G50, 60J15; secondary 60F16, 60F20.

Key words and phrases. Normed sums of independent random variables, integral tests.

THEOREM 1. [No assumption on $F(x)$]

- (i) $J_+(\gamma) = \infty$ iff $P\{\limsup S_n/\gamma_n = \infty\} = 1$.
- (ii) $J_-(\gamma) < J_+(\gamma) = \infty$ iff

$$P\{\liminf S_n/\gamma_n = \liminf(|X_1| + \cdots + |X_n|)/\gamma_n\}$$

$$= P\{\limsup S_n/\gamma_n = \infty\} = 1.$$
- (iii) $J_+(\gamma) + J_-(\gamma) < \infty$ iff $E\gamma^{-1}(|X|) < \infty$.

REMARK. Similar statements may be made, by symmetry, about the case where $J_-(\gamma) = \infty$ or the case where $J_+(\gamma) < J_-(\gamma) = \infty$. It follows from (1.1) that (iii) is implied by (i). However, (iii) is purely an analytic fact and a proof of (iii) is given in the next section by a simple nonprobabilistic argument.

Kesten (1970) and Erickson (1973) studied the case where $\gamma_n = n$ and obtained similar results. The integral test in Theorem 1 is the same as that in Theorem 2 of Erickson (1973) for $\gamma_n = n$. Like Kesten (1970), Theorem 1 is obtained by investigating the positive and negative contributions to the random walk $\{S_n\}$ with respect to the sequence γ_n . In fact, we have the following stronger

THEOREM 2. Suppose that either $E|X| = \infty$ or $\lim \gamma_n/n = \infty$. Then, one of the following alternatives must prevail:

- (i) $J_+(\gamma) = \infty$ and $P\{\limsup X_n^+ / (\gamma_n + X_1^- + \cdots + X_n^-) = \infty\} = 1$;
- (ii) $J_+(\gamma) < \infty$ and

$$P\{\lim(X_1^+ + \cdots + X_n^+) / (\gamma_n + X_1^- + \cdots + X_n^-) = 0\} = 1.$$

Theorem 1 and Theorem 2 will be proved in Section 2. We have two corollaries below which follow easily from Theorem 2.

COROLLARY 1. Suppose that $E|X| + \lim \gamma_n/n = \infty$. Then, $J_+(\gamma) < \infty$ iff

$$S_n/\gamma_n = \left(- \sum_{i=1}^n X_i^- / \gamma_n \right) (1 + o(1)) + o(1) \quad \text{a.s.}$$

COROLLARY 2. Suppose that $\lim \gamma_n/n = \infty$. Then, it is impossible for any random walk to have

$$(1.2) \quad -\infty < \liminf S_n/\gamma_n < 0 \quad \text{a.s.}$$

Erickson (1976, page 818) pointed out that it is impossible to have

$$0 < \liminf n^{-\alpha} S_n < \infty \quad \text{a.s. for } \alpha < 1.$$

Actually, it follows from his argument that it is impossible for any random walk to have

$$(1.3) \quad 0 < \liminf S_n/b_n < \infty \quad \text{a.s. for } b_n > 0 \text{ and } \lim b_n/n = 0.$$

Corollary 2 is the analogue of (1.3) for $\lim \gamma_n/n = \infty$.

2. Proofs. Lemma 1 gives an inequality for truncated expectations of partial sums of i.i.d. nonnegative random variables. The inequality has its own interest and may be used for other purposes.

LEMMA 1. Let $Y, Y_1, \dots, Y_n$ be i.i.d. nonnegative random variables. Set $S_n = Y_1 + \dots + Y_n$ and $m(x) = \int_0^x P\{Y \geq y\} dy$. Let $C > 0$ be a constant. Then

$$E \min(S_n, C) \leq \min(C, nm(C)) \leq 16E \min(S_n, C).$$

PROOF. Let $S'_n = \sum_{i=1}^n \min(Y_i, C)$. We shall discuss two cases, $nm(C) > 3C$ and $nm(C) \leq 3C$. For the first case, $nm(C) = uC > 3C$,

$$\begin{aligned} P\{S_n \leq C\} &= P\{ES'_n - S'_n \geq (u-1)C\} \\ &\leq (u-1)^{-2} C^{-2} nE(\min(Y, C))^2 \leq u/(u-1)^2 \leq 3/4, \end{aligned}$$

$$E \min(S_n, C) \geq CP\{S_n \geq C\} \geq C/4,$$

$$C \leq 16E \min(S_n, C).$$

For $nm(C) = uC \leq 3C$,

$$\begin{aligned} nm(C) &= ES'_n \\ &\leq 8E \min(S_n, C) + ES'_n I\{S'_n > 8C\}, \\ ES'_n I\{S'_n > 8C\} &\leq (nE(\min(Y, C))^2 + (nm(C))^2) / (8C) \\ &\leq (Cnm(C) + 3Cnm(C)) / (8C) = nm(C)/2, \\ nm(C) &\leq 16E \min(S_n, C). \end{aligned}$$

The inequality in the other direction is obvious. $\square$

We shall study the ratio of two independent nonnegative random walks instead of the ratio of the positive and negative contributions of the random walk $\{S_n, n \geq 1\}$.

THEOREM 3. Let $\{W_n\}$ and $\{V_n\}$ be two independent sequences of i.i.d. nonnegative random variables. Suppose that $EW_1 + EV_1 + \lim \gamma_n/n = \infty$. Then the following statements are equivalent:

- (i) $\lim(W_1 + \dots + W_n)/(\gamma_n + V_1 + \dots + V_n) = 0$ a.s.;
- (ii) $\limsup(W_1 + \dots + W_n)/(\gamma_n + V_1 + \dots + V_n) < \infty$ a.s.;
- (iii) $\limsup W_n/(\gamma_n + V_1 + \dots + V_n) < \infty$ a.s.;
- (iv) $\sum_{n=1}^{\infty} P\{\delta W_n > \gamma_n + V_1 + \dots + V_n\} < \infty$ for some $\delta > 0$;
- (v) $\sum_{n=1}^{\infty} P\{\delta W_n > \gamma_n + V_1 + \dots + V_n\} < \infty$ for all $\delta > 0$;
- (vi) $\int_0^{\infty} \min(\gamma^{-1}(x), x/\int_0^x P\{V_1 \geq t\} dt) dP\{W_1 \leq x\} < \infty$.

PROOF. It suffices to prove (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v) $\Rightarrow$ (vi) $\Rightarrow$ (iv) $\Rightarrow$ (i) since it is obvious that (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii).

(iii) $\Rightarrow$ (iv). There are constants $\delta > 0$ and $k < \infty$ such that

$$P\left\{\bigcup_{n=k}^{\infty} [\delta W_n > \gamma_n + V_1 + \dots + V_n]\right\} \leq 1 - \delta < 1.$$

Set

$$A_n = [\delta W_n > \gamma_n + V_1 + \cdots + V_n],$$

$$A_n A_{n+m} \subset A_n [\delta W_{n+m} > \gamma_m + V_{n+1} + \cdots + V_{n+m}],$$

$$P\left\{A_n \bigcup_{j=1}^{\infty} A_{n+jk}\right\} \leq P\{A_n\} P\left\{\bigcup_{j=1}^{\infty} A_{jk}\right\} \leq P\{A_n\}(1 - \delta),$$

$$\delta \sum_{n=1}^N P\{A_{nk+i}\} \leq \sum_{n=1}^N P\left\{A_{nk+i} \left[\bigcup_{j=1}^{\infty} A_{(n+j)k+i}\right]^c\right\} \leq 1, \quad i = 1, \dots, k.$$

(iv) $\Rightarrow$ (v). Since $\gamma(2n) \geq 2\gamma(n)$, we have

$$\begin{aligned} & \sum_{n=1}^{\infty} P\{2\delta W_n > \gamma_n + V_1 + \cdots + V_n\} \\ & \leq 1 + 2 \sum_{n=1}^{\infty} P\{2\delta W_{2n} > \gamma(2n) + V_1 + \cdots + V_{2n}\} \\ & \leq 1 + 4 \sum_{n=1}^{\infty} P\{\delta W_n > \gamma_n + V_1 + \cdots + V_n\}. \end{aligned}$$

(v) $\Rightarrow$ (vi) $\Rightarrow$ (iv). Let $T = \inf\{k: S_k > C\}$, $S_n = \sum_{i=1}^n V_i$, and $S'_k = \sum_{i=1}^k \min(V_i, C)$. Then, it follows from Wald's lemma that

$$\int_0^C P\{V_1 \geq t\} dt E \min(T, n) = ES'_{\min(T, n)},$$

$$\min(S_n, C) \leq S'_{\min(T, n)} \leq 2 \min(S_n, C),$$

$$E \min(T, n) = 1 + \sum_{k=1}^{n-1} P\{V_1 + \cdots + V_k \leq C\}.$$

Therefore, by Lemma 1,

$$\begin{aligned} & \sum_{n=1}^{\infty} P\{W_n > \gamma_n + V_1 + \cdots + V_n\} \\ & \leq \int_0^{\infty} \sum_{\gamma(1) \leq \gamma(n) < x} P\{V_1 + \cdots + V_n < x\} dP\{W_1 \leq x\} \\ & \leq 2 \int_0^{\infty} \min\left(\gamma^{-1}(x), x \int_0^x P\{V_1 \geq t\} dt\right) dP\{W_1 \leq x\} \\ & \leq 32 + 32 \int_0^{\infty} \sum_{\gamma(1) \leq \gamma(n) < x} P\{V_1 + \cdots + V_n \leq x\} dP\{W_1 \leq x\} \\ & \leq 32 + 32 \int_0^{\infty} \sum_{n=1}^{\infty} P\{V_1 + \cdots + V_n + \gamma_n < 2x\} dP\{W_1 \leq x\} \\ & = 32 + 32 \sum_{n=1}^{\infty} P\{2W_1 > \gamma_n + V_1 + \cdots + V_n\}. \end{aligned}$$

(iv) $\Rightarrow$ (i). We shall prove (v) $\Rightarrow$ (i) instead.

$$\begin{aligned} & \sum_{n=1}^{\infty} P\{\delta(W_1 + W_2) > \gamma_n + V_1 + \cdots + V_n\} \\ & \leq 1 + 2 \sum_{n=1}^{\infty} P\{\delta(W_1 + W_2) > \gamma(2n) + V_1 + \cdots + V_{2n}\} \\ & \leq 1 + 2 \sum_{n=1}^{\infty} P\{\delta W_1 > \gamma_n + V_1 + \cdots + V_n\} \\ & \quad + 2 \sum_{n=1}^{\infty} P\{\delta W_2 > \gamma_n + V_1 + \cdots + V_n\} \\ & < \infty. \end{aligned}$$

Let $T(M) = \inf\{n: \gamma_n + V_1 + \cdots + V_n + M \geq \delta(W_1 + W_2)\}$. Then, by (v), $ET(M) \leq ET(0) < \infty$ and we can choose large M such that $ET(M) < 2$. Set

$$T_1 = T^{(1)} = T(M),$$

$$T^{(n)} = \inf\{j: \gamma_j + V_{k+1} + \cdots + V_{k+j} + M \geq \delta(W_{2n-1} + W_{2n})\}$$

on $\{T_{n-1} = k\}$, and

$$T_n = T_{n-1} + T^{(n)}, \quad n = 2, 3, \dots$$

Then, by the strong law of large numbers, there exists an integer-valued random variable N such that

$$T_n = T^{(1)} + \cdots + T^{(n)} < 2n - 1 \quad \text{for any } n \geq N.$$

Since γ_n/n is nondecreasing,

$$\gamma(T^{(1)}) + \cdots + \gamma(T^{(n)}) \leq \gamma(T_n) < \gamma(2n - 1) \quad \text{for any } n \geq N,$$

$$\delta(W_1 + \cdots + W_{2n}) \leq nM + \gamma(2n - 1) + V_1 + \cdots + V_{2n-1}, \quad n \geq N.$$

By the condition that $EW_1 + EV_1 + \lim \gamma_n/n = \infty$,

$$nM = (\gamma_n + V_1 + \cdots + V_n + W_1 + \cdots + W_n)o(1) \quad \text{a.s.}$$

Hence,

$$\limsup(W_1 + \cdots + W_n)/(\gamma_n + V_1 + \cdots + V_n) \leq 1/\delta \quad \text{a.s.}$$

And (i) follows because that δ is arbitrary. $\square$

PROOF OF THEOREM 2. We may assume that $P\{X > 0\} \neq 0$ and $P\{X < 0\} \neq 0$. Let

$$T = T_1 = T^{(1)} = \inf\{k: X_k > 0\}, \quad T_0 = S_0 = 0,$$

$$T^{(n)} = \inf\{k > 0: X_{k+T_{n-1}} > 0\}, \quad T_n = T_{n-1} + T^{(n)}, \quad n \geq 2,$$

and

$$W_n = X_{T_n}, \quad V_n = -(S_{T_n} - S_{T_{n-1}} - W_n), \quad n \geq 1.$$

Since $T^{(n)}$ are copies of T , (W_n, V_n) , $n \geq 1$, are i.i.d. random vectors [see Chow and Teicher (1978, page 136, Lemma 3)]. It is not difficult to find that W_1 is

independent of V_1 [see also Kesten (1970, middle of page 1185)].

$$(2.1) \quad P\{W_1 > t\} = P\{X > t\} / P\{X > 0\}, \quad EW_1 \geq EX^+,$$

$$V_1 = \sum_{n=1}^T X_n^-, \quad EV_1 \geq EX^-,$$

$$(2.2) \quad \int_0^x P\{V_1 \geq t\} dt = E \min(V_1, x) \leq E \sum_{n=1}^T \min(X_n^-, x) \\ = ETE \min(X^-, x),$$

$$P\{T > n\} = (P\{X \leq 0\})^n,$$

$$ET < \infty \quad \text{and} \quad X_1^- \leq V_1.$$

Therefore, by (2.1) and (2.2),

$$(2.3) \quad J_+(\gamma) < \infty$$

iff

$$\int_0^\infty \min\left(\gamma^{-1}(x), x \int_0^x P\{V_1 \geq t\} dt\right) dP\{W_1 \leq x\} < \infty.$$

Let k be an integer with $ET < k$. By the strong law of large numbers.

$$P\{T_n \geq kn, \text{i.o.}\} = 0,$$

$$\begin{aligned} \limsup X_n^+ / (\gamma_n + X_1^- + \cdots + X_n^-) \\ &\geq \limsup W_n / (\gamma(T_n) + V_1 + \cdots + V_n) \\ &\geq \limsup W_n / (\gamma(kn) + V_1 + \cdots + V_{kn}) \\ &\geq \limsup W_{kn+i} / (\gamma(kn+i) + V_1 + \cdots + V_{kn+i}) \quad \text{for any } i \\ &= \limsup W_n / (\gamma_n + V_1 + \cdots + V_n) \quad \text{a.s.} \end{aligned}$$

It follows from Theorem 3(iii), (vi), and (2.3) that

$$(2.4) \quad P\{\limsup X_n^+ / (\gamma_n + X_1^- + \cdots + X_n^-) = \infty\} = 1 \quad \text{if } J_+(\gamma) = \infty.$$

Similarly,

$$\begin{aligned} \limsup (X_1^+ + \cdots + X_n^+) / (\gamma_n + X_1^- + \cdots + X_n^-) \\ &\leq \limsup (W_1 + \cdots + W_n) / (\gamma(T_{n-1}) + V_1 + \cdots + V_{n-1}) \\ &\leq \limsup (W_2 + \cdots + W_n) / (\gamma_{n-1} + V_2 + \cdots + V_n) \\ &= \limsup (W_1 + \cdots + W_n) / (\gamma_n + V_1 + \cdots + V_n) \quad \text{a.s.} \end{aligned}$$

And it follows from Theorem 3(i), (vi), and (2.3) that

$$P\{\limsup (X_1^+ + \cdots + X_n^+) / (\gamma_n + X_1^- + \cdots + X_n^-) = 0\} = 1,$$

if $J_+(\gamma) < \infty$. This completes the proof.

PROOF OF THEOREM 1. (i) and (ii) follow easily from Theorem 2 and Corollary 1. Since $J_+(\gamma) + J_-(\gamma) \leq E\gamma^{-1}(|X|)$, we only need to prove the only if part of (iii). Suppose that

$$(2.5) \quad J_+(\gamma) + J_-(\gamma) < \infty.$$

Since $E\gamma^{-1}(|X|) \leq \gamma^{-1}(1)[E|X| + 1]$, we can further assume that $E|X| = \infty$. Set

$$(2.6) \quad h(x) = \min\left(\gamma^{-1}(x), x / \int_0^x P\{|X| \geq t\} dt\right).$$

Then,

$$(2.7) \quad \begin{aligned} & \int_0^\infty h(x) dP\{|X| \leq x\} \\ &= \int_0^\infty \min\left(\gamma^{-1}(x), x / (m_+(x) + m_-(x))\right) dP\{|X| \leq x\} \\ &\leq J_+(\gamma) + J_-(\gamma) < \infty \quad \text{by (2.5).} \end{aligned}$$

Since $\gamma(x)/x$ is nondecreasing,

$$(2.8) \quad h(x) \text{ is nondecreasing and } h(x)/x \text{ is nonincreasing.}$$

Choose x_0 so that $\int_{x_0}^\infty h(x) dP\{|X| \leq x\} \leq 1/2$. Then by (2.8)

$$\begin{aligned} \int_0^y P\{|X| \geq t\} dt &= \int_0^\infty \min(t, y) dP\{|X| \leq t\} \\ &\leq x_0 + y(h(y))^{-1} \int_{x_0}^\infty h(t) dP\{|X| \leq t\}. \end{aligned}$$

Therefore, since $E|X| = \infty$,

$$h(y)y^{-1} \int_0^y P\{|X| \geq t\} dt \leq x_0 h(y)/y + 1/2 \rightarrow 1/2$$

and

$$h(y) = \gamma^{-1}(y) \quad \text{for all large } y.$$

This and (2.7) finish the proof.

Acknowledgment. The result of Corollary 2 was pointed out by the referee and he also simplified our proof of the present form of the paragraph after (2.8). His efforts and contributions are truly appreciated by us.

REFERENCES

- CHOW, Y. S. and TEICHER, H. (1978). *Probability Theory*. Springer, New York.
 DERMAN, C. and ROBBINS, H. (1955). The strong law of large numbers when the first moment does not exist. *Proc. Nat. Acad. Sci. U.S.A.* **41** 586–587.
 ERICKSON, K. B. (1973). The strong law of large numbers when the mean is undefined. *Trans. Amer. Math. Soc.* **185** 371–381.
 ERICKSON, K. B. (1976). Recurrence sets of normed random walk in R^d . *Ann. Probab.* **4** 802–828.
 FELLER, W. (1946). A limit theorem for random variables with infinite moments. *Amer. J. Math.* **68** 257–262.
 KESTEN, H. (1970). The limit points of a normalized random walk. *Ann. Math. Statist.* **41** 1173–1205.

DEPARTMENT OF STATISTICS
COLUMBIA UNIVERSITY
NEW YORK, NEW YORK 10027

DEPARTMENT OF APPLIED MATHEMATICS
AND STATISTICS
STATE UNIVERSITY OF NEW YORK
AT STONY BROOK
STONY BROOK, NEW YORK 11794