

Bivariate Tail Probability Approximations

John E. Kolassa and Donghyun Lee
Rutgers, the State University of New Jersey

21 December 2023

Motivation Example

- 1 Hao and Kolassa (2016) analyze a crossover study

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma
 - ② Measures efficacy of canine therapy on PTSD, vs standard therapy.

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma
 - ② Measures efficacy of canine therapy on PTSD, vs standard therapy.
 - ③ Performance was evaluated on both therapies for subject j : $X_j^i = 1$ if canine therapy superior for subject j with ordering $i \in \{1, 2\}$.

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma
 - ② Measures efficacy of canine therapy on PTSD, vs standard therapy.
 - ③ Performance was evaluated on both therapies for subject j : $X_j^i = 1$ if canine therapy superior for subject j with ordering $i \in \{1, 2\}$.
- ② Hypotheses:

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma
 - ② Measures efficacy of canine therapy on PTSD, vs standard therapy.
 - ③ Performance was evaluated on both therapies for subject j : $X_j^i = 1$ if canine therapy superior for subject j with ordering $i \in \{1, 2\}$.
- ② Hypotheses:
 - ① Null hypothesis is that there is no systematic difference in therapies, regardless of ordering.

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma
 - ② Measures efficacy of canine therapy on PTSD, vs standard therapy.
 - ③ Performance was evaluated on both therapies for subject j : $X_j^i = 1$ if canine therapy superior for subject j with ordering $i \in \{1, 2\}$.
- ② Hypotheses:
 - ① Null hypothesis is that there is no systematic difference in therapies, regardless of ordering.
 - ② Alternative hypothesis is that canine therapy is superior for at least one of the ordering.

Motivation Example

- ① Hao and Kolassa (2016) analyze a crossover study
 - ① Studies US military subject to physical trauma
 - ② Measures efficacy of canine therapy on PTSD, vs standard therapy.
 - ③ Performance was evaluated on both therapies for subject j : $X_j^i = 1$ if canine therapy superior for subject j with ordering $i \in \{1, 2\}$.
- ② Hypotheses:
 - ① Null hypothesis is that there is no systematic difference in therapies, regardless of ordering.
 - ② Alternative hypothesis is that canine therapy is superior for at least one of the ordering.
- ③ p -value is given by $P [\bar{X}^1 \geq \bar{x}^1, \bar{X}^2 \geq \bar{x}^2]$.

Notation

- 1 Random vector $\mathbf{X}_i = (X_i^1, \dots, X_i^p)$ has a continuous distribution,

Notation

- ① Random vector $\mathbf{X}_i = (X_i^1, \dots, X_i^p)$ has a continuous distribution,
- ② Moment generating function

$$M(\tau_1, \dots, \tau_p) = \mathbb{E} [\exp(\tau_1 X_i^1 + \dots + \tau_p X_i^p)] ,$$

Notation

- ① Random vector $\mathbf{X}_i = (X_i^1, \dots, X_i^p)$ has a continuous distribution,
- ② Moment generating function

$$M(\tau_1, \dots, \tau_p) = \mathbb{E} [\exp(\tau_1 X_i^1 + \dots + \tau_p X_i^p)] ,$$

- ① finite for $|\tau_i| < \epsilon$ for all $i \leq p$ and some $\epsilon > 0$.

Notation

- ① Random vector $\mathbf{X}_i = (X_i^1, \dots, X_i^p)$ has a continuous distribution,
- ② Moment generating function

$$M(\tau_1, \dots, \tau_p) = \mathbb{E} [\exp(\tau_1 X_i^1 + \dots + \tau_p X_i^p)] ,$$

- ① finite for $|\tau_i| < \epsilon$ for all $i \leq p$ and some $\epsilon > 0$.
- ③ Let the cumulant generating function be $K(\tau_1, \dots, \tau_p) = \log(M(\tau_1, \dots, \tau_p))$.

Notation

- ① Random vector $\mathbf{X}_i = (X_i^1, \dots, X_i^p)$ has a continuous distribution,
- ② Moment generating function

$$M(\tau_1, \dots, \tau_p) = \mathbb{E} [\exp(\tau_1 X_i^1 + \dots + \tau_p X_i^p)] ,$$

- ① finite for $|\tau_i| < \epsilon$ for all $i \leq p$ and some $\epsilon > 0$.
- ③ Let the cumulant generating function be $K(\tau_1, \dots, \tau_p) = \log(M(\tau_1, \dots, \tau_p))$.
- ④ Let $\hat{\tau}_j$ satisfy $K^k(\tau_1, \dots, \tau_p) = \bar{x}^k$ for $k = 1, \dots, p$.

Notation

- ① Random vector $\mathbf{X}_i = (X_i^1, \dots, X_i^p)$ has a continuous distribution,
- ② Moment generating function

$$M(\tau_1, \dots, \tau_p) = \mathbb{E} [\exp(\tau_1 X_i^1 + \dots + \tau_p X_i^p)] ,$$

- ① finite for $|\tau_i| < \epsilon$ for all $i \leq p$ and some $\epsilon > 0$.
- ③ Let the cumulant generating function be $K(\tau_1, \dots, \tau_p) = \log(M(\tau_1, \dots, \tau_p))$.
- ④ Let $\hat{\tau}_j$ satisfy $K^k(\tau_1, \dots, \tau_p) = \bar{x}^k$ for $k = 1, \dots, p$.
- ⑤ Let $\bar{\mathbf{X}} = \sum_{i=1}^n \mathbf{X}_i / n$

Objective: Approximate joint tail probabilities for \bar{X}

- 1 of n independent and identically distributed random vectors (X_i^1, \dots, X_i^p) .

Objective: Approximate joint tail probabilities for \bar{X}

- 1 of n independent and identically distributed random vectors (X_i^1, \dots, X_i^p) .
- 2 Use existence of higher moments to obtain better performance than CLT.

Objective: Approximate joint tail probabilities for \bar{X}

- 1 of n independent and identically distributed random vectors (X_i^1, \dots, X_i^p) .
- 2 Use existence of higher moments to obtain better performance than CLT.
 - 1 Gives Edgeworth series.

Objective: Approximate joint tail probabilities for \bar{X}

- ① of n independent and identically distributed random vectors (X_i^1, \dots, X_i^p) .
- ② Use existence of higher moments to obtain better performance than CLT.
 - ① Gives Edgeworth series.
- ③ Use existence of cumulant generating function to obtain better relative error behavior than Edgeworth series.

Objective: Approximate joint tail probabilities for \bar{X}

- ① of n independent and identically distributed random vectors (X_i^1, \dots, X_i^p) .
- ② Use existence of higher moments to obtain better performance than CLT.
 - ① Gives Edgeworth series.
- ③ Use existence of cumulant generating function to obtain better relative error behavior than Edgeworth series.
 - ① Measure theoretical error behavior in terms of inverse powers of \sqrt{n} .

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

① $\oint = \int_{c_1 - i\infty}^{c_1 + i\infty} \cdots \int_{c_p - i\infty}^{c_p + i\infty}$

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

① $\oint = \int_{c_1 - i\infty}^{c_1 + i\infty} \cdots \int_{c_p - i\infty}^{c_p + i\infty}$

- ② $\mathbf{c} = (c_1, \dots, c_p)$ in the interior of the domain of K .

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

① $\oint = \int_{c_1 - i\infty}^{c_1 + i\infty} \cdots \int_{c_p - i\infty}^{c_p + i\infty}$

- ② $\mathbf{c} = (c_1, \dots, c_p)$ in the interior of the domain of K .

- ② The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}] = P[\bar{X}^1 \geq \bar{x}^1 \cap \dots \cap \bar{X}^p \geq \bar{x}^p]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

① $\oint = \int_{c_1 - i\infty}^{c_1 + i\infty} \cdots \int_{c_p - i\infty}^{c_p + i\infty}$

- ② $\mathbf{c} = (c_1, \dots, c_p)$ in the interior of the domain of K .

- ② The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}] = P[\bar{X}^1 \geq \bar{x}^1 \cap \dots \cap \bar{X}^p \geq \bar{x}^p]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ① $c_i > 0$ for all i

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

① $\oint = \int_{c_1-i\infty}^{c_1+i\infty} \cdots \int_{c_p-i\infty}^{c_p+i\infty}$

- ② $\mathbf{c} = (c_1, \dots, c_p)$ in the interior of the domain of K .

- ② The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}] = P[\bar{X}^1 \geq \bar{x}^1 \cap \dots \cap \bar{X}^p \geq \bar{x}^p]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ① $c_i > 0$ for all i

- ③ When K is exactly quadratic, get Gaussian tail probabilities, exactly.

Inversion Integrals:

- ① Density $f_n(\bar{x}^1, \dots, \bar{x}^p)$ is

$$\frac{n^p}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) d\tau_1 \cdots d\tau_p$$

① $\oint = \int_{c_1 - i\infty}^{c_1 + i\infty} \cdots \int_{c_p - i\infty}^{c_p + i\infty}$

- ② $\mathbf{c} = (c_1, \dots, c_p)$ in the interior of the domain of K .

- ② The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}] = P[\bar{X}^1 \geq \bar{x}^1 \cap \dots \cap \bar{X}^p \geq \bar{x}^p]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ① $c_i > 0$ for all i

- ③ When K is exactly quadratic, get Gaussian tail probabilities, exactly.

- ① Write $\bar{\Phi}(\mathbf{z}) = P[Z^1 \geq z^1 \cap \dots \cap Z^p \geq z^p]$ for Z_i independent standard normals.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.
- 3 Exponentiate cubic and quartic terms.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.
- 3 Exponentiate cubic and quartic terms.
- 4 Integrate term-wise.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.
- 3 Exponentiate cubic and quartic terms.
- 4 Integrate term-wise.
- 5 Then $P[\bar{X}^1 \geq \bar{x}^1]$ is approximately

$$\exp(n[\hat{z}^2 - \hat{\omega}^2]) \left[\bar{\Phi}(\sqrt{n}\hat{z}) \left(1 - n\frac{\hat{\rho}_3}{6} \right) + \phi(\sqrt{n}\hat{z}) \frac{\hat{\rho}^3(n\hat{z}^2 - 1)}{6\sqrt{n}} \right].$$

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.
- 3 Exponentiate cubic and quartic terms.
- 4 Integrate term-wise.
- 5 Then $P[\bar{X}^1 \geq \bar{x}^1]$ is approximately

$$\exp(n[\hat{z}^2 - \hat{\omega}^2]) \left[\bar{\Phi}(\sqrt{n}\hat{z}) \left(1 - n\frac{\hat{\rho}_3}{6}\right) + \phi(\sqrt{n}\hat{z}) \frac{\hat{\rho}^3(n\hat{z}^2 - 1)}{6\sqrt{n}} \right].$$

- 1 $\hat{z} = \hat{\tau}\sqrt{K''(\hat{\tau})}$

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.
- 3 Exponentiate cubic and quartic terms.
- 4 Integrate term-wise.
- 5 Then $P[\bar{X}^1 \geq \bar{x}^1]$ is approximately

$$\exp(n[\hat{z}^2 - \hat{\omega}^2]) \left[\bar{\Phi}(\sqrt{n}\hat{z}) \left(1 - n\frac{\hat{\rho}_3}{6}\right) + \phi(\sqrt{n}\hat{z}) \frac{\hat{\rho}^3(n\hat{z}^2 - 1)}{6\sqrt{n}} \right].$$

- 1 $\hat{z} = \hat{\tau}\sqrt{K''(\hat{\tau})}$
- 2 $\hat{\omega} = \text{sign}(\hat{\tau})\sqrt{2(\hat{\tau}\bar{x} - K(\hat{\tau}))}$

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Expand $K(\tau) - \tau\bar{x}$ about $\hat{\tau}$:

$$K(\hat{\tau}) - \hat{\tau}\bar{x} + K''(\hat{\tau})(\tau - \hat{\tau})^2/2 + K'''(\hat{\tau})(\hat{\tau} - \hat{\tau})^3/6 + K''''(\tau^?)(\hat{\tau} - \hat{\tau})^4/24.$$

- 2 Set $c_1 = \hat{\tau}$.
- 3 Exponentiate cubic and quartic terms.
- 4 Integrate term-wise.
- 5 Then $P[\bar{X}^1 \geq \bar{x}^1]$ is approximately

$$\exp(n[\hat{z}^2 - \hat{\omega}^2]) \left[\bar{\Phi}(\sqrt{n}\hat{z}) \left(1 - n\frac{\hat{\rho}_3}{6}\right) + \phi(\sqrt{n}\hat{z}) \frac{\hat{\rho}_3(n\hat{z}^2 - 1)}{6\sqrt{n}} \right].$$

- 1 $\hat{z} = \hat{\tau}\sqrt{K''(\hat{\tau})}$
- 2 $\hat{\omega} = \text{sign}(\hat{\tau})\sqrt{2(\hat{\tau}\bar{x} - K(\hat{\tau}))}$
- 3 $\hat{\rho}_3 = K'''(\hat{\tau})K''(\hat{\tau})^{-3/2}$.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Re-parameterize the inversion integral in terms of ω satisfying $(\omega - \hat{\omega})^2/2 = K(\tau) - \tau x - K(\hat{\tau}) + \hat{\tau}x$.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Re-parameterize the inversion integral in terms of ω satisfying $(\omega - \hat{\omega})^2/2 = K(\tau) - \tau x - K(\hat{\tau}) + \hat{\tau}x$.
- 2 Integral is $\frac{1}{(2\pi i)} \oint \exp(n[\omega^2/2 - \hat{\omega}\omega]) \lambda \frac{d\omega}{\omega}$.

One Dimensional Methods: Hao and Kolassa (2016) approach

- 1 Re-parameterize the inversion integral in terms of ω satisfying $(\omega - \hat{\omega})^2/2 = K(\tau) - \tau x - K(\hat{\tau}) + \hat{\tau}x$.
- 2 Integral is $\frac{1}{(2\pi i)} \oint \exp(n[\omega^2/2 - \hat{\omega}\omega]) \lambda \frac{d\omega}{\omega}$.
 - 1 $\lambda = \frac{\omega}{\tau} \frac{d\tau}{d\omega}$.

One Dimensional Methods: Hao and Kolassa (2016) approach

- ① Re-parameterize the inversion integral in terms of ω satisfying $(\omega - \hat{\omega})^2/2 = K(\tau) - \tau x - K(\hat{\tau}) + \hat{\tau} x$.
- ② Integral is $\frac{1}{(2\pi i)} \oint \exp(n[\omega^2/2 - \hat{\omega}\omega]) \lambda \frac{d\omega}{\omega}$.
 - ① $\lambda = \frac{\omega}{\tau} \frac{d\tau}{d\omega}$.
- ③ Tail probability approximation is $\bar{\Phi}(\hat{\omega}) + \phi(\hat{\omega})(1/\hat{\omega} - 1/\hat{z})/\sqrt{n}$.

One Dimensional Methods: Hao and Kolassa (2016) approach

- ① Re-parameterize the inversion integral in terms of ω satisfying $(\omega - \hat{\omega})^2/2 = K(\tau) - \tau x - K(\hat{\tau}) + \hat{\tau} x$.
- ② Integral is $\frac{1}{(2\pi i)} \oint \exp(n[\omega^2/2 - \hat{\omega}\omega]) \lambda \frac{d\omega}{\omega}$.
 - ① $\lambda = \frac{\omega}{\tau} \frac{d\tau}{d\omega}$.
- ③ Tail probability approximation is $\bar{\Phi}(\hat{\omega}) + \phi(\hat{\omega})(1/\hat{\omega} - 1/\hat{z})/\sqrt{n}$.
 - ① Holds without regards to $\hat{\tau} > 0$.

Methods in Multiple Dimensions

- ① The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

Methods in Multiple Dimensions

- ① The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ② Hao and Kolassa (2016) works analogously as with Robinson:

Methods in Multiple Dimensions

- ① The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ② Hao and Kolassa (2016) works analogously as with Robinson:

- ① Expand $K(\tau) - \sum_j \tau_j \bar{x}^j$ about $\hat{\tau} = (\hat{\tau}_1, \dots, \hat{\tau}_p)$ to get:

$$\begin{aligned} K(\hat{\tau}) - \sum_j \hat{\tau}_j \bar{x}^j + \sum_{j,k} K^{jk}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)/2 \\ + \sum_{j,k,\ell} K^{jk\ell}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)/6 \\ + \sum_{j,k,\ell,m} K^{jk\ell m}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)(\tau_m - \hat{\tau}_m)/24. \end{aligned}$$

Methods in Multiple Dimensions

- ① The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ② Hao and Kolassa (2016) works analogously as with Robinson:

- ① Expand $K(\tau) - \sum_j \tau_j \bar{x}^j$ about $\hat{\tau} = (\hat{\tau}_1, \dots, \hat{\tau}_p)$ to get:

$$\begin{aligned} K(\hat{\tau}) - \sum_j \hat{\tau}_j \bar{x}^j + \sum_{j,k} K^{jk}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)/2 \\ + \sum_{j,k,\ell} K^{jk\ell}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)/6 \\ + \sum_{j,k,\ell,m} K^{jk\ell m}(\tau^?)(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)(\tau_m - \hat{\tau}_m)/24. \end{aligned}$$

- ② Expand exp of cubic and quartic terms.

Methods in Multiple Dimensions

- ① The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}_i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ② Hao and Kolassa (2016) works analogously as with Robinson:

- ① Expand $K(\tau) - \sum_j \tau_j \bar{x}_j$ about $\hat{\tau} = (\hat{\tau}_1, \dots, \hat{\tau}_p)$ to get:

$$\begin{aligned} K(\hat{\tau}) - \sum_j \hat{\tau}_j \bar{x}_j + \sum_{j,k} K^{jk}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)/2 \\ + \sum_{j,k,\ell} K^{jk\ell}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)/6 \\ + \sum_{j,k,\ell,m} K^{jk\ell m}(\tau^?)(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)(\tau_m - \hat{\tau}_m)/24. \end{aligned}$$

- ② Expand exp of cubic and quartic terms.
③ Integral of resulting quartic term is $O(1/n)$.

Methods in Multiple Dimensions

- ① The tail probability $P[\bar{\mathbf{X}} \geq \bar{\mathbf{x}}]$ is

$$\frac{1}{(2\pi i)^p} \oint \exp(n[K(\tau_1, \dots, \tau_p) - \sum_{i=1}^p \bar{x}^i \tau_i]) \frac{d\tau_1 \cdots d\tau_p}{\tau_1 \cdots \tau_p}.$$

- ② Hao and Kolassa (2016) works analogously as with Robinson:

- ① Expand $K(\tau) - \sum_j \tau_j \bar{x}^j$ about $\hat{\tau} = (\hat{\tau}_1, \dots, \hat{\tau}_p)$ to get:

$$\begin{aligned} K(\hat{\tau}) - \sum_j \hat{\tau}_j \bar{x}^j + \sum_{j,k} K^{jk}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)/2 \\ + \sum_{j,k,\ell} K^{jk\ell}(\hat{\tau})(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)/6 \\ + \sum_{j,k,\ell,m} K^{jk\ell m}(\tau^?)(\tau_j - \hat{\tau}_j)(\tau_k - \hat{\tau}_k)(\tau_\ell - \hat{\tau}_\ell)(\tau_m - \hat{\tau}_m)/24. \end{aligned}$$

- ② Expand exp of cubic and quartic terms.
③ Integral of resulting quartic term is $O(1/n)$.
④ Integrate terms up to cubic term-wise.

Downsides of existing approximation

- 1 Lots of terms

Downsides of existing approximation

- ① Lots of terms
- ② Terms are not particularly interpretable.

Downsides of existing approximation

- ① Lots of terms
- ② Terms are not particularly interpretable.
 - ① (Multivariate) normal tail probability is multiplied by an exponential factor.

Downsides of existing approximation

- ① Lots of terms
- ② Terms are not particularly interpretable.
 - ① (Multivariate) normal tail probability is multiplied by an exponential factor.
- ③ You are stuck with $\hat{\tau}_j > 0$

Downsides of existing approximation

- ① Lots of terms
- ② Terms are not particularly interpretable.
 - ① (Multivariate) normal tail probability is multiplied by an exponential factor.
- ③ You are stuck with $\hat{\tau}_j > 0$
 - ① Get around this using Boole's Law

Lugannani and Rice Analog

- 1 New parameterization $\sum_j \hat{\omega}_j^2/2 = K(\hat{\tau}) - \sum_j \hat{\tau}_j$,
 $\sum_j (\omega_j - \hat{\omega}_j)^2/2 = K(\tau) - \sum_j \tau_j - (K(\hat{\tau}) - \sum_j \hat{\tau}_j)$

Lugannani and Rice Analog

- 1 New parameterization $\sum_j \hat{\omega}_j^2/2 = K(\hat{\tau}) - \sum_j \hat{\tau}_j$,
 $\sum_j (\omega_j - \hat{\omega}_j)^2/2 = K(\tau) - \sum_j \tau_j - (K(\hat{\tau}) - \sum_j \hat{\tau}_j)$
- 2 Integral is

$$\frac{1}{(2\pi i)^p} \oint \exp\left(n \sum_j \omega_j^2 - \sum_{i=1}^p \hat{\omega}_j \omega_j\right) \lambda \frac{d\omega_1 \cdots d\omega_p}{\omega_1 \cdots \omega_p}.$$

Lugannani and Rice Analog

- 1 New parameterization $\sum_j \hat{\omega}_j^2/2 = K(\hat{\tau}) - \sum_j \hat{\tau}_j$,
 $\sum_j (\omega_j - \hat{\omega}_j)^2/2 = K(\tau) - \sum_j \tau_j - (K(\hat{\tau}) - \sum_j \hat{\tau}_j)$
- 2 Integral is

$$\frac{1}{(2\pi i)^p} \oint \exp(n \sum_j \omega_j^2 - \sum_{i=1}^p \hat{\omega}_j \omega_j) \lambda \frac{d\omega_1 \cdots d\omega_p}{\omega_1 \cdots \omega_p}.$$

- 1 $\lambda = \frac{\omega_1 \cdots \omega_p}{\tau_1 \cdots \tau_p} \frac{d\tau_1 \cdots d\tau_p}{d\omega_1 \cdots d\omega_p}.$

Lugannani and Rice Analog

- ① New parameterization $\sum_j \hat{\omega}_j^2/2 = K(\hat{\tau}) - \sum_j \hat{\tau}_j$,
 $\sum_j (\omega_j - \hat{\omega}_j)^2/2 = K(\tau) - \sum_j \tau_j - (K(\hat{\tau}) - \sum_j \hat{\tau}_j)$
- ② Integral is

$$\frac{1}{(2\pi i)^p} \oint \exp\left(n \sum_j \omega_j^2 - \sum_{i=1}^p \hat{\omega}_j \omega_j\right) \lambda \frac{d\omega_1 \cdots d\omega_p}{\omega_1 \cdots \omega_p}.$$

① $\lambda = \frac{\omega_1 \cdots \omega_p}{\tau_1 \cdots \tau_p} \frac{d\tau_1 \cdots d\tau_p}{d\omega_1 \cdots d\omega_p}.$

- ① In one dimension, zeros in denominator of λ correspond to zeros in the numerator.

Lugannani and Rice Analog

- ① New parameterization $\sum_j \hat{\omega}_j^2/2 = K(\hat{\tau}) - \sum_j \hat{\tau}_j$,
 $\sum_j (\omega_j - \hat{\omega}_j)^2/2 = K(\tau) - \sum_j \tau_j - (K(\hat{\tau}) - \sum_j \hat{\tau}_j)$
- ② Integral is

$$\frac{1}{(2\pi i)^p} \oint \exp\left(n \sum_j \omega_j^2 - \sum_{i=1}^p \hat{\omega}_j \omega_j\right) \lambda \frac{d\omega_1 \cdots d\omega_p}{\omega_1 \cdots \omega_p}.$$

① $\lambda = \frac{\omega_1 \cdots \omega_p}{\tau_1 \cdots \tau_p} \frac{d\tau_1 \cdots d\tau_p}{d\omega_1 \cdots d\omega_p}.$

- ① In one dimension, zeros in denominator of λ correspond to zeros in the numerator.
- ② This fails for $p > 1$.

Re-parameterize to fix lack of alignment of zeros

- 1 Re-parameterize to $\xi_j = \omega_j - \sum_{k < j} \omega_k \zeta_j^k(\omega_1, \dots, \omega_{j-1})$ so that $\xi_j = 0$ if and only if $\tau_j = 0$.

Re-parameterize to fix lack of alignment of zeros

- 1 Re-parameterize to $\xi_j = \omega_j - \sum_{k < j} \omega_k \zeta_j^k(\omega_1, \dots, \omega_{j-1})$ so that $\xi_j = 0$ if and only if $\tau_j = 0$.
 - 1 Ex. $\xi_1 = \omega_1$, $\xi_2 = \omega_2 - \omega_1 \zeta_2^1(\omega_1)$ for $\zeta_1^2 = \omega_2(\tau_1, 0)/\omega_1$.

Re-parameterize to fix lack of alignment of zeros


- ① Re-parameterize to $\xi_j = \omega_j - \sum_{k < j} \omega_k \zeta_j^k(\omega_1, \dots, \omega_{j-1})$ so that $\xi_j = 0$ if and only if $\tau_j = 0$.
 - ① Ex. $\xi_1 = \omega_1$, $\xi_2 = \omega_2 - \omega_1 \zeta_2^1(\omega_1)$ for $\zeta_2^1 = \omega_2(\tau_1, 0)/\omega_1$.
 - ② Adjustment also makes $\{\bar{X}^1 \geq \bar{x}^1 \cap \dots, \bar{X}^p \geq \bar{x}^p\} \approx \{\hat{\Xi}^1 \geq \hat{\xi}^1 \cap \dots, \hat{\Xi}^p \geq \hat{\xi}^p\}$.

Re-parameterize to fix lack of alignment of zeros

- ① Re-parameterize to $\xi_j = \omega_j - \sum_{k < j} \omega_k \zeta_j^k(\omega_1, \dots, \omega_{j-1})$ so that $\xi_j = 0$ if and only if $\tau_j = 0$.
 - ① Ex. $\xi_1 = \omega_1$, $\xi_2 = \omega_2 - \omega_1 \zeta_2^1(\omega_1)$ for $\zeta_2^1 = \omega_2(\tau_1, 0)/\omega_1$.
 - ② Adjustment also makes $\{\bar{X}^1 \geq \bar{x}^1 \cap \dots, \bar{X}^p \geq \bar{x}^p\} \approx \{\hat{\Xi}^1 \geq \hat{\xi}^1 \cap \dots, \hat{\Xi}^p \geq \hat{\xi}^p\}$.
 - ③ Adjustment makes singularities in λ removable.

Adjustment induces sample space rotation for $\hat{\Omega}_j$


① using $\zeta_k^j(\hat{\Omega})$.



Correlated gamma example, $\mathbf{x} = (4, 5)$

Adjustment induces sample space rotation for $\hat{\Omega}_j$


① using $\zeta_k^j(\hat{\Omega})$.



Correlated gamma example, $\mathbf{x} = (4, 5)$

Adjustment induces sample space rotation for $\hat{\Omega}_j$

① using $\zeta_k^j(\hat{\Omega})$.



Correlated gamma example, $\mathbf{x} = (4, 5)$

New approximation is simpler

$$\textcircled{1} \quad \mathbb{P} \left[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p \right] \approx \\ \bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j) (1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$$

New approximation is simpler

- ① $P[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p] \approx$
 $\bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j)(1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$
① Σ is generated by transformation from ω to ξ .

New approximation is simpler

- ① $P[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p] \approx$
 $\bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j)(1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$
- ① Σ is generated by transformation from ω to ξ .
 - ② σ_j is a standard error for $\hat{\tau}_j$.

New approximation is simpler

- ① $P[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p] \approx$
 $\bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j)(1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$
 - ① Σ is generated by transformation from ω to ξ .
 - ② σ_j is a standard error for $\hat{\tau}_j$.
- ② In order to make λ have removable singularities,

New approximation is simpler

- ① $P[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p] \approx$
 $\bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j)(1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$
 - ① Σ is generated by transformation from ω to ξ .
 - ② σ_j is a standard error for $\hat{\tau}_j$.
- ② In order to make λ have removable singularities,
 - ① exponent in inversion integral is no longer exactly quadratic.

New approximation is simpler

- ① $P[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p] \approx$
 $\bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j)(1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$
 - ① Σ is generated by transformation from ω to ξ .
 - ② σ_j is a standard error for $\hat{\tau}_j$.
- ② In order to make λ have removable singularities,
 - ① exponent in inversion integral is no longer exactly quadratic.
 - ② C adjusts for this.

New approximation is simpler

- ① $P[X^1 \geq x^1 \cap \dots \cap X^p \geq x^p] \approx$
 $\bar{\Phi}(\sqrt{n}\hat{\xi}; \Sigma) + \sum_{j=1}^p \bar{\Phi}(\sqrt{n}\hat{\xi}_{-j}; \Sigma_j) \phi(\sqrt{n}\xi_j)(1/\xi_j - 1/(\hat{\tau}_j/\sigma_j)) + C.$
 - ① Σ is generated by transformation from ω to ξ .
 - ② σ_j is a standard error for $\hat{\tau}_j$.
- ② In order to make λ have removable singularities,
 - ① exponent in inversion integral is no longer exactly quadratic.
 - ② C adjusts for this.
- ③ Avoids requirement that $\hat{\tau}_j > 0$.

Example

① $p = 2,$


Example

- ① $p = 2$,
- ② $X_1 = Z_1 + Z_2$, $X_2 = Z_1 + Z_3$, Z_j independent exponentials.


Example

- ① $p = 2$,
- ② $X_1 = Z_1 + Z_2$, $X_2 = Z_1 + Z_3$, Z_j independent exponentials.
- ③ $n = 1!$


Relative Error: Normal on raw scale




Relative Error: Normal on SRLLR scale



Relative Error: SRLLR scale, adjust for marginal non-normality



Relative Error: SRLLR scale, all adjustments



Work in various stages of completion

- 1 Addressing removable singularities in C .

Work in various stages of completion

- 1 Addressing removable singularities in C .
- 2 $p > 2$.

Work in various stages of completion

- 1 Addressing removable singularities in C .
- 2 $p > 2$.
- 3 r^* version placing $O(1/\sqrt{n})$ corrections into argument of $\bar{\Phi}$.

Work in various stages of completion

- 1 Addressing removable singularities in C .
- 2 $p > 2$.
- 3 r^* version placing $O(1/\sqrt{n})$ corrections into argument of $\bar{\Phi}$.
- 4 Extension to lattice variables

Work in various stages of completion

- 1 Addressing removable singularities in C .
- 2 $p > 2$.
- 3 r^* version placing $O(1/\sqrt{n})$ corrections into argument of $\bar{\Phi}$.
- 4 Extension to lattice variables
- 5 Extension to conditional distributions.

Hao, Y. and Kolassa, J. (2016). Multiple hypotheses in the analysis of a crossover trial. *Statistics in Biosciences*, 8(2):253–263. Funding Information: The second author was supported by Grant NSF DMS 0906569. Publisher Copyright: © 2016, International Chinese Statistical Association.

Hao, Y. and Kolassa, J. (2016). Multiple hypotheses in the analysis of a crossover trial. *Statistics in Biosciences*, 8(2):253–263. Funding Information: The second author was supported by Grant NSF DMS 0906569. Publisher Copyright: © 2016, International Chinese Statistical Association.

Kolassa, J. (2003). Multivariate saddlepoint tail probability approximations. *Annals of Statistics*, 31:274–286.

Hao, Y. and Kolassa, J. (2016). Multiple hypotheses in the analysis of a crossover trial. *Statistics in Biosciences*, 8(2):253–263. Funding Information: The second author was supported by Grant NSF DMS 0906569. Publisher Copyright: © 2016, International Chinese Statistical Association.

Kolassa, J. (2003). Multivariate saddlepoint tail probability approximations. *Annals of Statistics*, 31:274–286.

Lugannani, R. and Rice, S. (1980). Saddle point approximation for the distribution of the sum of independent random variables. *Advances in Applied Probability*, 12:475–490.

- Hao, Y. and Kolassa, J. (2016). Multiple hypotheses in the analysis of a crossover trial. *Statistics in Biosciences*, 8(2):253–263. Funding Information: The second author was supported by Grant NSF DMS 0906569. Publisher Copyright: © 2016, International Chinese Statistical Association.
- Kolassa, J. (2003). Multivariate saddlepoint tail probability approximations. *Annals of Statistics*, 31:274–286.
- Lugannani, R. and Rice, S. (1980). Saddle point approximation for the distribution of the sum of independent random variables. *Advances in Applied Probability*, 12:475–490.
- Robinson, J. (1982). Saddlepoint approximations for permutation tests and confidence intervals. *Journal of the Royal Statistical Society Series B*, 44:91–101.